

Hanle effect in Coherent Backscattering

G. Labeyrie^{*†}, C. Miniatura^{*}, C. A. Müller^{*}, O. Sigwarth[†], D. Delande[†] and R. Kaiser^{*}

^{*} *Laboratoire Ondes et Désordre, FRE 2302 CNRS, 1361 route des Lucioles, F-06560 Valbonne.*

[†] *Laboratoire Kastler Brossel, Université Pierre et Marie Curie, 4 Place Jussieu, F-75005 Paris.*

⁺ *e-mail : labeyrie@inln.cnrs.fr*

(February 2, 2008)

We study the shape of the coherent backscattering (CBS) cone obtained when resonant light illuminates a thick cloud of laser-cooled rubidium atoms in presence of a homogenous magnetic field. We observe new magnetic field-dependent anisotropies in the CBS signal. We show that the observed behavior is due to the modification of the atomic radiation pattern by the magnetic field (Hanle effect in the excited state). PACS numbers: 42.20-y, 32.80.Pj

When a multiply scattering medium is illuminated by a laser beam, the scattered intensity results from the interference between the amplitudes associated with the various scattering paths; for a disordered medium, the interference terms are washed out when averaged over many sample configurations, *except* in a narrow angular range around exact backscattering where the average intensity is enhanced. This phenomenon, known as coherent backscattering (CBS), is due to a two-wave constructive interference (at exact back-scattering) between waves following a given scattering path and the associated reverse path, where exactly the same scatterers are visited in the reversed order [1]. The maximum enhancement is obtained when the amplitudes of the interfering paths are exactly balanced. For a convenient choice of polarization, time-reversal symmetry directly implies the equality of the interfering amplitudes. More generally, the interference phenomenon is very robust and qualitatively insensitive to most characteristics of the sample and illuminating wave.

However, applying a magnetic field on the sample breaks the time-reversal invariance. It was predicted [2], then experimentally observed [3] and theoretically studied [4,5] that it results in a decrease of the CBS enhancement as well as some rather complicated behavior of the cone shape. In our current understanding of CBS, two ingredients are essential : the individual scattering event, characterized for example by the radiation pattern of each scatterer (this is the single scattering ingredient), and the propagation in the medium between scattering events (this is the "average effective medium" ingredient). Of course, these two ingredients are not independent since the optical theorem links the propagation in the medium to the individual scattering. In the presence of a magnetic field, both ingredients can be affected. For the propagation, this is well known under the name of Faraday effect (and the Voigt or Cotton-Mouton effect) and can be described by the modification of the complex refractive index by the magnetic field. The magnetic

field-induced variation of the radiation pattern is much less studied, because it is very small in usual magneto-optically active materials; it is responsible for the "photonic Hall effect" predicted by van Tiggelen [6] and later observed experimentally [7]. In this paper, we show a novel situation where it is experimentally and theoretically possible to discriminate between the two ingredients and where the modification of the radiation pattern dominates the propagation effects.

In our experiment, we analyze coherent backscattering of resonant light by a dilute gas of laser-cooled rubidium atoms. Some features of this medium differ markedly from those used in previous CBS experiments. First, the cold atomic cloud constitutes a monodisperse sample of point scatterers, highly resonant in the vicinity of the optical transition. This implies a dramatic increase of the scattering cross section at resonance, but also a great sensitivity to any external perturbation like a magnetic field. Typically, few Gauss are enough to bring the atom completely off resonance, in sharp contrast with previous studies where Teslas were needed to induce significant effects [3]. Concurrently, a giant Faraday effect is observed in the cold atomic cloud [8]. Another important difference with classical systems is the presence of a quantum internal structure, which has been shown to strongly reduce the coherent backscattering interference [9]. Obviously, the addition of a magnetic field will affect the contrast of the CBS cone in various ways, and we have indeed observed new behaviors that will be described somewhere else. In the present Letter, we focus on the shape of the cone rather than its intensity.

The experiment has been described elsewhere [9]. We use the $J = 3 \rightarrow J' = 4$ transition of the D2 line of Rb^{85} (wavelength $\lambda = 780$ nm, natural width $\Gamma/2\pi = 5.9$ MHz) where J here denotes the hyperfine angular momentum. We show in Fig. 1 some CBS cones obtained in the linear polarization channels, for different values of the magnetic field and laser detuning. After scaling and subtraction of the incoherent background (average intensity in the wings of the peak), we plot the detected far-field intensity isolines. The horizontal and vertical coordinates correspond to the two azimuthal scattering angles. The magnetic field $\mathbf{B}$ is orthogonal to the plane of the figure, like the backscattered light wavevector $\mathbf{k}_b$, which points towards the reader. The incident light polarization is parallel to the vertical axis of the figure. Since we will care about the sign of magneto-optical ro-

FIG. 1. Experimental CBS cones. The far-field backscattered intensity is plotted as a function of the two azimuthal scattering angles (maximum intensity at center; we use a logarithmic scale for the isolines for a better visualization of the wings where the anisotropy is more pronounced; the lowest isoline corresponds to roughly 10% of the peak value). The total angular range is 2 mrad. $\mathbf{P}$ denotes the incident polarization. (a) $B = 0$, top: $\text{lin} \parallel \text{lin}$, bottom: $\text{lin} \perp \text{lin}$; (b) both cones in $\text{lin} \perp \text{lin}$, top: $B = -8\text{G}$, bottom: $B = +8\text{G}$; note the 90° flip when B changes sign. (c) $B = +10\text{G}$ and $\delta_L = -2.6\Gamma$, top: $\text{lin} \parallel \text{lin}$, bottom: $\text{lin} \perp \text{lin}$.

For resonant light (laser detuning $\delta_L = \omega_L - \omega_A = 0$ where ω_L and ω_A are the laser and atomic frequencies respectively) and zero magnetic field $B = 0$, we obtain in the $\text{lin} \perp \text{lin}$ channel a cushion-shaped cone with a four-fold symmetry shown in Fig. 1(a) (bottom). In the parallel channel (top), the cone has an elliptical shape with its large axis parallel to the incident polarization. In the presence of a magnetic field, we observe in the $\text{lin} \perp \text{lin}$ channel (b) that the cone has now an elliptical shape with an inclination of 45° from the incident polarization, which does not depend on the magnetic field strength. It is remarkable that, when the sign of the magnetic field is reversed, the cone flips by 90° . In the $\text{lin} \parallel \text{lin}$ channel, the shape of the cone does not significantly change with the magnetic field. When the laser is detuned with respect to the zero-field atomic transition, the cones in both the $\text{lin} \parallel \text{lin}$ and $\text{lin} \perp \text{lin}$ channels become isotropic, as shown in Fig. 1(c).

To understand the features observed in Fig. 1, we need to remember that the CBS cone shape is closely linked to the distribution of distances between first and last visited scatterers in the sample. In fact, it is the Fourier transform of the transverse intensity profile of the scattered light when the sample is illuminated by a narrow light

beam [3]. Thus, any scattering asymmetry inside the medium immediately results in a cone shape asymmetry. For the discussion of the observed effect, it turns out (see below) that the internal structure (Zeeman sublevels) of the atom is not crucial. We thus first consider the simplest case where the atomic transition is $J = 0 \rightarrow J' = 1$. The Zeeman effect induces a splitting of the atomic resonance line in 3 components separated by μB where B is the magnetic field and $\mu/2\pi = 1.4 \text{ MHz/G}$ is the Zeeman shift rate. As soon as the Zeeman shift is comparable to the resonance width (magnetic field of the order of 4 Gauss), the scattering cross-section of the atom is strongly modified. For simplicity, we restrict the discussion to a resonant excitation $\delta_L = 0$. In this case, the response of the atom to a linearly polarized field is that of a dipole rotated from the incident polarization by an angle $\varphi = \arctan(2\mu B/\Gamma)$, in the plane perpendicular to B . This rotation is responsible for the well-known fluorescence dip curve observed in typical Hanle experiments in the excited state [11]. The effect of the magnetic field on the average effective medium is also well known: in the direction of the magnetic field, it induces the Faraday effect, i.e. a rotation of the electric field around the magnetic field. For a weak magnetic field $\mu B \ll \Gamma$, the Faraday angle θ_F per mean-free path is: $\theta_F \simeq -\mu B/\Gamma$ [8]. It is crucial to note that it has a sign *opposite* to that of the atomic dipole rotation. As CBS involves both propagation and individual scattering, the experimental observation of the sign of the rotation makes it possible to determine the dominant effect. In our case, we will show that the rotation of the atomic dipole leads.

We propose a simple model which explains the observed cone shape behavior, taking into account only the rotation of the atomic dipole. We will restrict the discussion to the case of double scattering, as it is known to be dominant in our experimental conditions and because anisotropy effects are expected to decrease for higher order scattering. Due to the exponential attenuation of light inside the sample, the first and last scatterers of most paths lie in a "skin layer" of thickness approximately one scattering mean-free path under the surface of the sample. Thus, we will assume that all propagation takes place in the transverse plane, orthogonal to the incident wave vector. In addition, we will neglect all propagation effects (i.e. Faraday and Cotton-Mouton).

Let us now consider an initial polarization vector $\mathbf{P}$. The first scatterer radiates a wave at an angle Ψ from $\mathbf{P}$ towards the second atom, which then radiates in the backscattering direction where the field is analyzed along the polarization $\mathbf{A}$ at an angle α with $\mathbf{P}$. The angular dependence of the interference term between reverse paths is readily given by $\sin^2(\varphi - \Psi) \sin^2(\varphi + \Psi - \alpha)$. This formula (valid for resonant light) allows us to understand most of the observed behavior of the CBS interference pattern. Using a general symmetry argument, one can show that the CBS cone should remain symmetric

FIG. 2. Theoretical CBS cones (double scattering). (a) and (b) use the $J = 0 \rightarrow J' = 1$ transverse plane model described in text (same parameters as in Fig.1). (c): cones in $lin \parallel lin$ (top) and $lin \perp lin$ (bottom) for a $J = 3 \rightarrow J' = 4$ transition and $B = 0$ (semi-infinite medium); the cones shapes are rounded off compared to (a), but the symmetry properties are identical.

Figure 2 shows some examples of the double scattering CBS cones obtained in the linear channels with this model. (a) and (b) correspond to the same parameters as in Fig. 1. One clearly sees ((a), bottom row) the

distinctive clover-leaf shape in the $lin \perp lin$ channel at $B = 0$ with 4 lobes along the bisectors (at 45°). At non-vanishing magnetic field one of the lobe pairs is favored depending on the sign of φ (b). The comparison between Fig. 1 and 2 confirms that the CBS cone tilt observed experimentally is consistent with the sign of the dipole rotation, and that our simple model catches the essential part of the physics. For off-resonant excitation $\delta_L \neq 0$, the picture becomes more complex since the induced dipole is elliptical. However, if the Zeeman splitting is large enough $\mu B \gg \Gamma$ and if the laser frequency is tuned to resonance with one of the circular components $\delta_L = \pm \mu B$, then the dipole is essentially circular and the cone is isotropic in agreement with the data in Fig. 1(c).

We now discuss the effect of the internal structure in the ground state, which is neglected in the previous model. Due to its internal structure, the rubidium atom does not behave like a pure dipole scatterer. More specifically, the radiation pattern depends on the ground state magnetic sublevels m_g and m'_g respectively before and after the scattering event. To determine the total CBS signal, one has to sum over all possible transitions ($m_{g1} \rightarrow m'_{g1}$, $m_{g2} \rightarrow m'_{g2}$) of two atoms. We have computed numerically the total CBS signal in this case and found that, for moderate B values (up to a few Gauss), the symmetry of the CBS pattern is only weakly modified compared to the $J = 0 \rightarrow J' = 1$ case. Indeed, in this regime, the CBS signal is dominated by the "Rayleigh" transitions $m_g \rightarrow m_g$, yielding the same behavior as in the $J = 0 \rightarrow J' = 1$ situation. Thus, and this is the important point, we can still use the results of the simple $J = 0 \rightarrow J' = 1$ model with an average dipole rotation φ to describe the cone behavior for the $J = 3 \rightarrow J' = 4$ transition of rubidium. One illustration of the effect of the internal structure is given in Fig. 2(c) where we plot the double scattering CBS signal at $B = 0$ in the $lin \parallel lin$ (top) and $lin \perp lin$ (bottom) channels for the $J = 3 \rightarrow J' = 4$ transition and a semi-infinite geometry. The main effect of the internal structure in this case is a rounding-off of the cone features. The comparison with the experimental cones of Fig. 1(a) yields a nice agreement. In order to make a quantitative test of our model, we have experimentally measured the dipole rotation φ (open squares in Fig. 3(a)) by analyzing the polarization of resonant light scattered by an optically-thin atomic cloud (optical thickness 0.05). In this regime, Faraday rotation and multiple scattering are negligible. The solid line in Fig. 3(a) corresponds to the theory for a $J = 3 \rightarrow J' = 4$ transition and agrees well with our measurements. In a different experiment, we have measured the Faraday rotation *per mean free path* θ_F (solid squares in Fig. 3(a)) by recording, as a function of B , the polarization rotation of light transmitted by a cloud of fixed optical thickness 1. When the magnetic field increases, the total scattering cross-section decreases yielding an increasing mean-free path. Thus, in order to keep the

factor in the presence of a magnetic field.

We thank the CNRS and the PACA Region for financial support. Laboratoire Kastler Brossel is laboratoire de l'Université Pierre et Marie Curie et de l'Ecole Normale Supérieure, UMR 8552 du CNRS.

FIG. 3. Magneto-optical rotations. (a): measured dipole rotation φ ($\square$) and Faraday rotation *per mean free path* θ_F ($\blacksquare$). The curves correspond to the theory for a $J = 3 \rightarrow J' = 4$ transition (see text for sign conventions). Note that the two rotations have opposite signs. (b): measured CBS flip angle α_f ($\blacktriangle$) compared to the $J = 3 \rightarrow J' = 4$ model neglecting all propagation effects (solid line).

In conclusion, we reported in this paper the observation of various anisotropies in the shape of the coherent backscattering cone from cold rubidium atoms in a magnetic field. The observed behavior differs radically from previous work on magneto-optically active samples where the cone features were determined by the Faraday effect during the propagation between scatterers. In our situation, the observed anisotropy is due to the modification of the atomic radiation pattern by the magnetic field. This observation illustrates the new regimes that are accessible through the use of cold atoms as a scattering medium for light. Some more elaborate modelling is needed to understand in detail the roles of propagation versus individual scattering effects in our sample, in particular to explain the behavior of the CBS enhancement

-
- [1] B. A. van Tiggelen, A. Lagendijk, and A. Tip, J. Phys. Condens. Matter **2**, 7653 (1990).
 - [2] A. A. Golubentsev, Zh. Eksp. Teor. Fiz. **86**, 47 (1984).
 - [3] F. A. Erbacher, R. Lenke and G. Maret, Europhys. Lett. **21**, 551 (1993); R. Lenke and G. Maret, Eur. Phys. J. B **17**, 171 (2000); R. Lenke, R. Lehner and G. Maret, Europhys. Lett. **52**, 620 (2000).
 - [4] F. C. MacIntosh and S. John, Phys. Rev. B **37**, 1884 (1988); A. S. Martinez and R. Maynard, Phys. Rev. B **50**, 3714 (1994); B. A. van Tiggelen, R. Maynard, and T. M. Nieuwenhuizen, Phys. Rev. E **53**, 2881 (1996).
 - [5] D. Lacoste and B. A. van Tiggelen, Phys. Rev. E **61**, 4556 (2000).
 - [6] B. A. van Tiggelen, Phys. Rev. Lett. **53**, 2881 (1996).
 - [7] G. L. J. A. Rikken and B. A. van Tiggelen, Nature **381**, 54 (1996).
 - [8] G. Labeyrie, C. Miniatura and R. Kaiser, Phys. Rev. A **64**, 033402 (2001).
 - [9] G. Labeyrie, F. de Tomasi, J.-C. Bernard, C. A. Müller, C. Miniatura and R. Kaiser, Phys. Rev. Lett. **83**, 5266 (1999); G. Labeyrie, C.A. Müller, D.S. Wiersma, C. Miniatura and R. Kaiser, J. Opt. B: Quantum Semi-class. Opt. **2**, 672 (2000); T. Jonckheere, C.A. Müller, R. Kaiser, C. Miniatura and D. Delande, Phys. Rev. Lett. **85**, 4269 (2000); C. A. Müller, T. Jonckheere, C. Miniatura and D. Delande, Phys. Rev. A **64**, 053804 (2001).
 - [10] M. P. van Albada, M. B. van der Mark, and A. Lagendijk, Phys. Rev. Lett. **58**, 361 (1987).
 - [11] W. Hanle, Z. Phys. **30**, 93 (1924); W. Rasmussen, R. Schieder, and H. Walter, Opt. Commun. **12**, 315 (1974).
 - [12] contrarily to what was stated in [8], where the dependence of the scattering mean-free path with magnetic field was overlooked.